

**§ 556.620**

**§ 556.620 Sulfabromomethazine sodium.**

Tolerances for residues of sulfabromomethazine sodium in food are established as follows:

(a) In the uncooked edible tissues of cattle at 0.1 part per million (negligible residue).

(b) In milk at 0.01 part per million (negligible residue).

[47 FR 30244, July 13, 1982]

**§ 556.625 Sodium sulfachloropyrazine monohydrate.**

A tolerance of zero is established for residues of sodium sulfachloropyrazine monohydrate in the uncooked edible tissues of chickens.

**§ 556.630 Sulfachlorpyridazine.**

A tolerance of 0.1 part per million is established for negligible residues of sulfachlorpyridazine in uncooked edible tissues of calves and swine.

**§ 556.640 Sulfadimethoxine.**

(a) [Reserved]

(b) *Tolerances.* (1) A tolerance of 0.1 part per million (ppm) is established for negligible residues of sulfadimethoxine in uncooked edible tissues of chickens, turkeys, cattle, ducks, salmonids, catfish, and chukar partridges.

(2) A tolerance of 0.01 ppm is established for negligible residues of sulfadimethoxine in milk.

[64 FR 26672, May 17, 1999]

**§ 556.650 Sulfaethoxypyridazine.**

Tolerances for residues of sulfaethoxypyridazine in food are established as follows:

(a) Zero in the uncooked edible tissues of swine and in milk.

(b) 0.1 part per million (negligible residue) in uncooked edible tissues of cattle.

**§ 556.660 Sulfamerazine.**

A tolerance of zero is established for residues of sulfamerazine (N<sup>1</sup>-[4-methyl-2-pyrimidinyl]sulfanilamide) in the uncooked edible tissues of trout.

**§ 556.670 Sulfamethazine.**

A tolerance of 0.1 part per million is established for negligible residues of

**21 CFR Ch. I (4–1–14 Edition)**

sulfamethazine in the uncooked edible tissues of chickens, turkeys, cattle, and swine.

[47 FR 25323, June 11, 1982]

**§ 556.685 Sulfaquinoxaline.**

A tolerance of 0.1 part per million is established for negligible residues of sulfaquinoxaline in the uncooked edible tissues of chickens, turkeys, calves, and cattle.

[61 FR 24443, May 15, 1996]

**§ 556.690 Sulfathiazole.**

A tolerance of 0.1 part per million is established for negligible residues of sulfathiazole in the uncooked edible tissues of swine.

**§ 556.700 Sulfomyxin.**

A tolerance of zero is established for residues of sulfomyxin (N-sulfomethyl-polymyxin B sodium salt) in uncooked edible tissues from chickens and turkeys.

**§ 556.710 Testosterone propionate.**

No residues of testosterone, resulting from the use of testosterone propionate, are permitted in excess of the following increments above the concentrations of testosterone naturally present in untreated animals:

(a) In uncooked edible tissues of heifers:

(1) 0.64 part per billion in muscle.

(2) 2.6 parts per billion in fat.

(3) 1.9 parts per billion in kidney.

(4) 1.3 parts per billion in liver.

(b) [Reserved]

[52 FR 27683, July 23, 1987]

**§ 556.720 Tetracycline.**

(a) *Acceptable daily intake (ADI).* The ADI for total tetracycline residues (chlortetracycline, oxytetracycline, and tetracycline) is 25 micrograms per kilogram of body weight per day.

(b) *Tolerances.* Tolerances are established for the sum of tetracycline residues in tissues of calves, swine, sheep, chickens, and turkeys, of 2 parts per million (ppm) in muscle, 6 ppm in liver, and 12 ppm in fat and kidney.

[63 FR 57246, Oct. 27, 1998]